

PRE AND POST CARE INSTRUCTIONSFORBOTOX

Pre Care Instructions

- Avoid the use of Aspirin, NSAIDS, Ginko Biloba, Garlic Supplements, Green Tea, Flax Oil,CodLiver Oil, Vitamins A and E, and essential fatty acids for up to 1 week pre and post injection as they will increase your risk ofbruising.
- Do not drink alcoholic beverages 24 hours before (or after) your treatment to avoid extra bruising
- You may take Arnica tablets 2-3 days prior to your injection to reduce the risk ofbruising
- Do not apply products that are potentially irritating for 2 -3 days before and after treatment (examples- tretinoin/retin-a, glycolic acid, benzoyl peroxide,hydroquinone)
- Inform your provider if you have a history of cold sores to receive a prescription for an antiviral prior totreatment

Day of Treatment

- Arrive with a clean face. Please do not wearmakeup.
- You may experience a mild amount of tenderness or stinging during and followinginjection
- Redness and swelling are normal. Some bruising may also bevisible

Post Care Instructions

- You can exercise your treated muscles for 1-2 hours after treatment (e.g. practice frowning, raise your eyebrows, and squint). This may help to work the product into yourmuscles.
- Stay in a vertical position for 4 hours following injection. Do not rest your head or lie down; sit upright.
- Ice (but avoid firm pressure) for 15 minutes 3 times the day of treatment ifneeded.
- You may take Acetaminophen/Tylenol if you experience any mildtenderness/discomfort
- Do not massage treated areas aftertreament
- After treatment there may be mild swelling and redness with possibility of bruising. You may apply or take Arnica tablets to help decrease the amount ofbruising
- Avoid extended UV exposure until any redness/swelling has subsided. ApplySPF.
- Avoid strenuous exercise for 24 hours to reduce bruising. You may resume othernormalactivities/routinesimmediately
- Avoid laser, IPL, or skin tightening treatments of the area for a minimum of 24 hours (or as directed by your provider) afterinjection
- You may continue your topical products as mentioned above 1 day after injection.